



Loki, Io: New ground-based observations and a model describing the change from periodic overturn

J. A. Rathbun¹ and J. R. Spencer²

Received 5 June 2006; revised 17 June 2006; accepted 24 June 2006; published 6 September 2006.

[1] Loki Patera is the most powerful volcano in the solar system. We have obtained measurements of Loki's 3.5 micron brightness from NASA's Infrared Telescope Facility (IRTF) and have witnessed a major change in eruptive behavior. While Loki brightened by a factor of several every 540 days prior to 2001, from 2001 through 2004 Loki remained at a constant, intermediate brightness. We have constructed a quantitative model of Loki as a basaltic lava lake whose solidified crust overturns when it becomes buoyantly unstable. By altering the speed at which the overturn propagates across the patera, we can match our ground-based brightness data. In addition, we can match other data taken at other times and wavelengths. By slowing the propagation speed dramatically, we can match the observations from 2001–2004. Such slowing could be due to a small change in volatile content in the lava. **Citation:** Rathbun, J. A., and J. R. Spencer (2006), Loki, Io: New ground-based observations and a model describing the change from periodic overturn, *Geophys. Res. Lett.*, 33, L17201, doi:10.1029/2006GL026844.

1. Introduction

[2] Loki Patera is a horseshoe-shaped dark feature 200 km across located on the Jupiter-facing hemisphere of the volcanically active moon Io. It is the site of prodigious amounts of thermal emission, at times accounting for nearly 15% of Io's total heat flow [Spencer *et al.*, 2000]. While its appearance at visible wavelengths has changed little since it was observed by the Voyager spacecraft [McEwen *et al.*, 1998; Geissler *et al.*, 2004], its near infrared brightness can vary by an order of magnitude over a timescale of several months [Rathbun *et al.*, 2002].

1.1. Telescopic Observations

[3] Loki's brightness in the infrared is large enough to be measured with ground-based telescopes using occultation techniques [Spencer *et al.*, 1990]. In this technique, Io is viewed while in eclipse, thereby eliminating reflected sunlight from the brightness measurement, and undergoing an occultation by Jupiter. Io's total brightness is measured photometrically as a function of time and each hotspot appears as a step in the resulting lightcurve. Loki's brightness has been measured in this manner for more than a decade [Rathbun *et al.*, 2002]. Its variability, indicative of fluctuations in the high-temperature component, is apparent.

Rathbun *et al.* [2002] tested the variability and found that the brightness from 1988 through 2001 varied periodically, with a period of 540 days.

[4] We have continued to measure Loki's brightness from NASA's Infrared Telescope Facility (IRTF) using the occultation technique (Figure 1). Although the temporal sampling rate has decreased, a change in Loki's behavior clearly occurred in mid-2001. Data taken before this time show two distinct populations of brightness: one low and one high. Data obtained between late 2001 and early 2003 show a single brightness population at a level between the previous "high" and "low" levels. In late 2003 the brightness fades to a low level and increases to a high level, indicating that another brightening event has begun. This new event appears to be out of phase with the earlier brightening events, but the temporal sampling has been inadequate to fully characterize Loki's recent behavior.

1.2. Spacecraft Data

[5] From 1999 through 2001 the Galileo spacecraft performed a series of close flybys of Io and obtained high resolution images from multiple instruments. Spencer *et al.* [2000] examined temperature measurements of Loki taken by the Galileo PhotoPolarimeter-Radiometer (PPR) in October 1999 and February 2000. Observations of Loki were taken at night to avoid reflected sunlight so the brightness measurements are dominated by low-temperature (≤ 300 K) emission. They found that the hottest part of Loki moved from the southwest corner in 1999 to the eastern edge in 2000. Images from October 1999 show the temperature gradually increasing away from the southwest corner. Approximately 70 km away, the temperature abruptly drops by more than 100 K. Rathbun *et al.* [2002] interpreted this as evidence of a front of hot, young material moving to the east. They further noted that Voyager images taken in March and July of 1979 also show evidence of a front of dark material moving to the east. Since an active plume was present near Loki at this time, they interpret this material to also be young.

[6] Rathbun *et al.* [2002] favor the interpretation that Loki is an overturning lava lake. In order to match these data, they suggest that a front of crustal foundering moves across Loki. As the crust cools and thickens, it eventually becomes negatively buoyant. One piece sinks, followed by the piece next to it and thus the resurfacing progresses in a coherent manner. So, the Galileo PPR data, which capture the low-temperature emission, show the warmer part of Loki that has already overturned and the cooler temperatures to the east where older material not yet overturned. Davies [2003] analyzed a high resolution infrared image from the Near Infrared Mapping Spectrometer (NIMS). Although the hot front was not seen in the image, the surface ages inferred from the observation varied nearly

¹Department of Physics, University of Redlands, Redlands, California, USA.

²Southwest Research Institute, Boulder, Colorado, USA.




Figure 1. The 3.5 μm brightness of Loki as measured primarily from Jupiter occultations. Some of the data was taken at other wavelengths (3.8, 4.8, and 3.39 μm); see *Rathbun et al.* [2002] for details. The dotted square wave has a period of 540 days to show the early periodicity.

linearly across Loki decreasing to the northeast, consistent with the overturn of a lava lake in the manner proposed by *Rathbun et al.* [2002].

2. Model

[7] Here, we quantify the model of *Rathbun et al.* [2002] and determine whether, in addition to matching the PPR data, it can quantitatively match the high-temperature emission that dominates the integrated 3.5 micron flux that is measured from the ground and if it can yield insight into the behavior changes observed in those data.

[8] Loki Patera has an area of approximately $2.1 \times 10^4 \text{ km}^2$ and a width of 55 km. For simplicity, we model Loki as a rectangular lava lake 390 km long and 55 km wide (as if the horseshoe of Loki were straightened; Figure 2). The depth of the lava lake is not set, but we assume it is deep enough that the bottom does not affect our calculations. We model the age of the surface of the lake as a function of position and time. The age is assumed constant along the short dimension. The initial age of the surface, created in the previous wave of foundering, varies linearly along the length of the Patera, with the oldest surface in the west. Based on the early ground-based observations [*Rathbun et al.*, 2002], we set the oldest crust at 540 days and the youngest at 150 days. As time progresses, crust on the west begins to overturn, with the overturn front moving to the east. The rate at which the overturn progresses is the only major variable in the model.

[9] Once the age of the surface is calculated as a function of distance and time, we use the lava cooling model of *Davies et al.* [2005] to calculate the temperature of the surface. This finite-element model assumes a basaltic composition of the lava with a liquidus temperature of 1200–1600 K. Finally, knowing the area and surface temperature of each raft and assuming blackbody emission, we determine the total brightness of Loki at 3.5 microns, simulating what would be seen from the ground (most ground-based data record only Loki’s integrated thermal emission).

[10] At spatial resolutions better than 100 m (5.5 km^2 in area), our model is independent of resolution. Since no rafts are observed in the Galileo data and calculation time for the model increases dramatically with higher resolution, we use a compromise spatial resolution of 2 m (area of 0.1 km^2) in our computations. Although we calculate the brightness of

Loki at intervals of 1 day, the surface ages can be fractions of a day. When a particular length of crust has overturned on a day, say x pixels, the age of this crust varies from 1 day to $1/x$ days.

[11] Altering the speed of the foundering front has two affects. Obviously, the faster the overturn propagates, the shorter the brightening event will last. Further, the faster it propagates, the more surface area younger than a given age is produced, thus increasing the total brightness. We find that the brightness is proportional to speed with the constant of proportionality approximately 32 when the brightness is measured in $\text{GW}/\mu\text{m}/\text{str}$ and the propagation speed in km/day .

3. Model Results

[12] The variations observed in the ground-based data (Figure 1) can be explained by variations in the overturn propagation speed. Approximately 9 brightening events can be seen in the data, but only five have well defined starting and ending times. The average duration of these 5 brightenings is 225 days, requiring a propagation speed of 1.7 km/day , equivalent to an areal production rate of $1080 \text{ m}^2/\text{s}$. Given this speed, the model predicts an average active brightness of $55 \text{ GW}/\mu\text{m}/\text{str}$, very close to the average active brightness of $60 \text{ GW}/\mu\text{m}/\text{str}$ observed during these 5 events (Figure 3b). Thus, our single variable model is able to match both the observable quantities: duration and brightness.

3.1. Sulfur

[13] Measurements of Loki’s thermal emission by Voyager’s Infrared Interferometer Spectrometer (IRIS) don’t require temperatures higher than 450 K [*Pearl and Sinton*, 1982]. This relatively low temperature led to the suggestion that Loki might be a sulfur lake [*Lunine and Stevenson*, 1985], though the IRIS data can also be fitted with silicate models [*Carr*, 1986; *Howell*, 1997]. For completeness, we checked whether our ground-based 3.5 micron data could be reconciled with a sulfur lake model.




Figure 2. A graphical depiction of our model. The top view shows how temperature varies along the 390 km length of the lava length. The side view shows how the crust thickens with time until eventually sinking below the liquid magma and a new crust begins to form.




Figure 3. Model results. (a) The solid line shows the modeled 3.5 μm brightness as a function of time for the period from May 1997 through September 2000. The diamonds are observed brightnesses. (b) Duration and average brightnesses possible from model for both basalt and sulfur. Plus signs show the duration and average brightness for each of the five observed brightening events. The asterisk shows the average duration and brightness for all observed events. (c) The locations of overturn front on July 12th and August 4th, 1998 (asterisks) based on our model. The black boxes show the position of the hottest material seen on those dates by *Macintosh et al.* [2003]. The horseshoe shows the long dimension of the model wrapped around the patera. Background image from *Macintosh et al.* [2003].

[14] While the *Davies et al.* [2005] cooling model for basalt is numerical, *Howell* [1997] constructs an analytic formulation in terms of the chemical properties of the magma. He finds that $T(t) = a_T t^{-1/8}$ where $a_T = \left(\frac{1}{\sqrt{\pi}} \left[\frac{\sqrt{K\rho C}}{\text{Erf}(\Lambda)} \right] \frac{\Delta T}{\sigma} \right)^{1/4}$, K is the thermal conductivity, ρ is the density, C is the heat capacity, σ is the Stefan-Boltzman constant, and Λ is the solution to the transcendental equation $\frac{L}{C\Delta T} = \frac{e^{-\Lambda^2}}{\Lambda\sqrt{\pi}\text{Erf}(\Lambda)}$ where L is the heat of fusion. Since sulfur is a single element, its material properties are better constrained than those of basalt. We use values of $K = 0.205$ W/mK, $\rho = 1.819$ g/cm³, $C = 22.75$ J/(mol K), $\Delta T = 100$ K, $L = 1.727$ kJ/mol, and an atomic mass of 32.065 g/mol to find that $a_T = 225.7$ K. Due to the resolution of our model, the maximum temperatures reached are well below the boiling point of sulfur. With this modification to our model, we again attempted to match the 3.5 micron brightness of Loki. We find that for a given front speed, the calculated brightness is nearly three orders of magnitude lower for sulfur compared to basalt (Figure 3b). The duration of the brightening event depends only on speed, not composition, so if we increase the modeled speed to match the observed bright-

ness, the duration will not match. From this, we conclude that Loki patera is not an overturning lake of liquid sulfur.

3.2. Modeling Individual Brightenings, 1997–2000

[15] Since our basalt model was able to match the average brightness and duration of a brightening event, we attempted to match the brightening changes witnessed during an event. The best temporal resolution data for Loki were taken during the period of 1997 through 2000, during the Galileo era. We began by changing velocity at times when the data showed a large (>10 GW/micron/str) change in brightness. While this matched the 1999–2000 brightening very well, for the earlier events we added some time segments when the velocity changed linearly with time. Remarkably, not only were we able to match the observed brightness, but the event also ends when observed (Figure 3a). The mean velocity for this modeled period is 1.53 km/day. Since the material overturning is 55 km wide, the rate at which new material is exposed at the surface is 970 m²/s, comparable to the rate of 1160 m²/s found by *Howell et al.* [2001]. *Howell* was specifically looking at October and November of 1999, where our velocities are 1.8 km/day, equivalent to an exposure rate of 1150 m²/s.

[16] Using speckle techniques, *Macintosh et al.* [2003] observed Io and were able to measure Loki's 2.2 micron brightness on July 12th and 28th and August 4th, 1998. We ran the same model from the 1997 thru 2001 period, but calculating the 2.2 micron brightness instead. The model again matches the observed brightnesses to within 10%. The nearest 3.5 micron observations were taken June 19th and August 20th, so a direct comparison of the model with multiple wavelengths simultaneously is not possible. *Macintosh et al.* [2003] were also able to determine the position of the Loki hot spot on July 12th and August 4th, 1998. Using the same velocity profile from the model, we calculated the location of the overturning front on those dates to be 64 and 86 km from the western edge, respectively. To compare these locations with *Macintosh's* observations we need to take our linear model and transform it back into a horseshoe shape. Figure 3c shows a white horseshoe representing the long dimension in our model with asterisks at the positions we calculated for the front on the two days. The background is an image from *Macintosh et al.* [2003] showing a Voyager image of Loki and two error boxes (in black) for the position of the hottest area observed on those dates. Our modeled positions fall within these error boxes.

3.3. Modeling Data From 2001

[17] In 2001, Loki's 3.5 micron behavior changed. No longer did the brightness alternate between "bright" and "dark". Instead, it remained at a reasonably constant 30–45 GW/micron/str for at least 500 and up to 900 days. A brightening event was observed to begin between September 15th and October 10th, 1999. The brightening was still occurring on March 7th, 2000. No data was taken between then and December 24th, 2000. In that time, we infer that brightening event ended. Between December 24th, 2000 and April 14th, 2002 fifteen observations were made with an average brightness of 37 GW/micron/str and a smaller than normal deviation from this average. Loki's brightness then remained near the "dark" level until a new "bright"




Figure 4. Results of model for 2001 through 2004. (a) Brightness of Loki in 2001 as measured by *Marchis et al.* [2005] at different wavelengths. The lines are modeled brightnesses at those wavelengths assuming a speed of 0.5 km/day. (b) Modeled brightness as a function of time assuming a wave speed of 0.5 km/day and a second front beginning 540 days after the first.

event began in late 2003, out of phase with the previous events.

[18] Simply reducing the velocity of the overturn propagation allows our model to match the brightnesses observed in 2001–2002. A speed of 0.9 km/day yields an event with a maximum brightness of 29 GW/micron/str and a duration of 450 days, a reasonable fit to observations. This velocity is similar to that calculated by *Davies* [2003]. He fit temperatures to high spatial resolution NIMS spectra of Loki taken October 10th, 2001. Those data did not show the front of hot material, likely because the observation covered only the southern part of the patera, but from the temperature variation across the Patera, he calculated a resurfacing rate of 1 km/day.

[19] Observations of Io using adaptive optics can isolate the brightness from individual volcanoes. *Marchis et al.* [2005] measured Loki’s brightness on December 18th, 20th, and 28th, 2001 at 2.2, 3.8, and 4.3 microns, near of middle of the intermediate brightness period. These data show much more variability at 3.8 and 4.3 microns than is seen in the occultation data. Because these data capture only a very small portion of a brightening event, we attempted only to match the average brightnesses measured (Figure 4a). While a speed of 0.5 km/day reasonably matches the observations, this is slightly lower than the average value of 0.9 km/day used to match the average 3.5 micron data, but the 1997–2000 data show that variations in velocity by a factor of 2 are not unusual.

[20] When velocities are as low as 0.5 km/day, it takes 780 days to overturn the entire patera. We had previously assumed that a new brightening even began after 540 days when the solidified crust was again of sufficient density to founder. In the past, this new overturn always began after the previous one was finished. Furthermore, simply by changing the velocity to 0.5 km/day, the easternmost rafts are already older than 540 days when they sink, so it is not clear that a when a new overturn will begin. However, we model the case of simultaneous overturn fronts for completeness. We assume both fronts have a velocity of 0.5 km/day, both fronts begin at the western edge, and that the second front begins 540 days after the first (Figure 4b). The maximum brightness occurs when both fronts are present and is 33 GW/micron/str, twice the 17 GW/micron/str when

only one front is active. So, another way to match the observations from 2001 through 2002 is for there to be simultaneous fronts for part of the time, perhaps at different speeds. Without more data, we cannot unambiguously determine which mechanism accounts for the change in Loki’s behavior, but a change in velocity is required in either case.

4. Discussion

[21] We have shown that a simple quantitative model of an overturning basaltic lava lake is consistent with observations by a variety of observers at a variety of wavelengths using a variety of instruments and techniques including ground-based occultation measurements at 3.5 microns, ground-based speckle measurements at 2.2 microns plus position, ground-based adaptive optics measurements at 2.2, 3.8, and 4.3 microns, and Galileo PPR and NIMS data at mid-infrared and far-infrared wavelengths. All the data can be matched by simple changes in the velocity of the overturn propagation. But, why does the velocity change?

[22] A change in velocity indicates a change in the age of the surface when it sinks. In the original *Rathbun et al.* [2002] model the rafts overturn when their density increases to the point where they are denser than the underlying liquid in the lake. In order for the solid crust to not sink immediately, there must be some porosity in the crust that lowers its density to lower than the liquid. Porosity measurements for a Hawaiian lava lake [*Peck et al.*, 1966] show




Figure 5. The density of the solidified crust as a function of time assuming the porosity profile of *Peck et al.* [1966].

surface porosities near 40%, decreasing nearly exponentially to about 10% at depth, similar to that calculated by *Matson et al.* [2006] for Loki considering SO₂ content and pressure in the lava lake. We use a simple Stefan model to calculate the thickness of the solidified crust as a function of time and incorporate the *Peck et al.* [1966] measurements to calculate the density of the solidified crust as a function of time (Figure 5). After 400 days, the density remains remarkably constant, with only a 1% variation between 400 and 800 days. This is due to the fact that the thickness changes little at this time and also the porosity changes little with depth, resulting in little variation of density at that time. This shows that small differences in density of the underlying lava will yield large differences in the age of the raft when it founders. Further, small differences in initial density (especially porosity) of the crust can also yield large differences in age. Finally, other factors, such as the behavior of neighboring slabs, can also have an effect on when a particular raft sinks. We believe the most likely explanation for the variations in age at which the raft sinks is small changes in lava volatile content, which affects both the lava and crust density. These small changes can produce large variations in the sinking age and thus the propagation speed of the sinking front.

[23] **Acknowledgments.** For some of this work, J. A. R. was a visiting astronomer at the Infrared Telescope Facility, which is operated by the University of Hawaii under Cooperative Agreement NCC 5-538 with the National Aeronautics and Space Administration, Science Mission Directorate, Planetary Astronomy Program. We thank Robert Howell and Laszlo Keszthelyi for very helpful reviews.

References

- Carr, M. H. (1986), Silicate volcanism on Io, *J. Geophys. Res.*, *91*, 3521–3532.
- Davies, A. G. (2003), Temperature, age and crust thickness distributions of Loki Patera on Io from Galileo NIMS data: Implications for resurfacing mechanism, *Geophys. Res. Lett.*, *30*(21), 2133, doi:10.1029/2003GL018371.
- Davies, A. G., D. L. Matson, G. J. Veeder, T. V. Johnson, and D. L. Blaney (2005), Post-solidification cooling and the age of Io's lava flows, *Icarus*, *176*, 123–137.
- Geissler, P., A. McEwen, C. Phillips, L. Keszthelyi, and J. Spencer (2004), Surface changes on Io during the Galileo mission, *Icarus*, *169*, 29–64.
- Howell, R. R. (1997), Thermal emission from lava flows on Io, *Icarus*, *127*, 394–407.
- Howell, R. R., et al. (2001), Ground-based observations of volcanism on Io in 1999 and early 2000, *J. Geophys. Res.*, *106*, 33,129–33,139.
- Lunine, J. L., and D. J. Stevenson (1985), Physics and chemistry of sulfur lakes on Io, *Icarus*, *64*, 345–367.
- Macintosh, B. A., D. Gavel, S. G. Gibbard, C. E. Max, M. Eckart, I. dePater, A. M. Ghez, and J. Spencer (2003), Speckle imaging of volcanic hotspots on Io with the Keck telescope, *Icarus*, *165*, 137–143.
- Marchis, F., et al. (2005), Keck AO survey of Io global volcanic activity between 2 and 5 μm , *Icarus*, *176*, 96–122.
- Matson, D. L., A. G. Davies, G. J. Veeder, J. A. Rathbun, T. V. Johnson, and J. C. Castillo (2006), Io: Loki Patera as a magma sea, *J. Geophys. Res.*, doi:10.1029/2006JE002703, in press.
- McEwen, A. S., et al. (1998), Active volcanism on Io as seen by Galileo SSI, *Icarus*, *135*, 181–219.
- Pearl, J. C., and W. M. Sinton (1982), Hot spots of Io, in *Satellites of Jupiter*, edited by D. Morrison, pp. 724–755, Univ. of Ariz. Press, Tucson.
- Peck, D. L., T. L. Wright, and J. G. Moore (1966), Crystallization of theleitic basalt in Alae lava lake, Hawaii, *Bull. Volcanol.*, *29*, 629–655.
- Rathbun, J. A., J. R. Spencer, A. G. Davies, R. R. Howell, and L. Wilson (2002), Loki, Io: A periodic volcano, *Geophys. Res. Lett.*, *29*(10), 1443, doi:10.1029/2002GL014747.
- Spencer, J. R., M. A. Shure, M. E. Ressler, J. D. Goguen, W. M. Sinton, D. W. Toomey, A. Denault, and J. Westfall (1990), Discovery of hotspots on Io using disk-resolved infrared imaging, *Nature*, *348*, 618–621.
- Spencer, J. R., J. A. Rathbun, L. D. Travis, L. K. Tamppari, L. Barnard, T. Z. Martin, and A. S. McEwen (2000), Io's thermal emission from the Galileo Photopolarimeter-Radiometer, *Science*, *288*, 1198–1201.
- J. A. Rathbun, Department of Physics, University of Redlands, 1200 E. Colton Ave., Redlands, CA 92374, USA. (julie_rathbun@redlands.edu)
- J. R. Spencer, Southwest Research Institute, 1050 Walnut St., Suite 400, Boulder, CO 80302, USA. (spencer@boulder.swri.edu)